

P - Lindelöf Spaces

Dr. Radhi I.M.Ali *

(Accepted 16/8/2005)

□ ABSTRACT □

In this work, we study p-Lindelöf spaces, a concept which is "stronger" than Lindelöf spaces. Every p-compact space is p-Lindelöf space, so the concept of p-Lindelöf is a generalization of the concept of p-compact. Most of the theorems which are valid for p-compact spaces will be also valid for p-Lindelöf spaces, (with a slight modification). We also study locally p-Lindelöf spaces.

*Assistant Prof. Dept of Math, Jerash Private Univ. Jerash - Jordan.

فضاءات p - لندلوف

الدكتور راضي إبراهيم محمد علي*

(قبل للنشر في 2005/8/16)

□ الملخص □

درسنا في هذا البحث فضاءات p - لندلوف التي هي "أقوى" من فضاءات لندلوف. لأن كل فضاء p - متراص هو p - لندلوف فإن مفهوم فضاء p - لندلوف هو تعميم لمفهوم p - متراص. إن معظم المبرهنات التي تصح في فضاءات p - متراصة تصح كذلك في فضاءات p - لندلوف مع استخدام بعض المعالجات الطفيفة. كما عرفنا الفضاءات التي تتصف بأنها p - لندلوف موضعياً (محلياً) وحصلنا على بعض النتائج المتعلقة بها.

* أستاذ مساعد في قسم الرياضيات - كلية الزراعة و العلوم - جامعة جرش الأهلية - جرش - الأردن.

1. Introduction:

In a recent paper [5], the authors introduce and study p -compact spaces. In this paper we study p -Lindelöf spaces as a generalization of p -compact space. Most of the theorems which are valid for p -compact will be valid for p -Lindelöf spaces. In order to make this work as self-contained as possible we are going to give some preliminaries theorems and definitions we need them later.

Definition 1.1 [1]: A topological space (X, τ) is Lindelöf space if every open cover of X has a countable subcover.

Definition 1.2 [5]: A subset A of a topological space (X, τ) is called pre-open if $A \subseteq (A)^\circ$. A subset B is a pre-closed if $X \setminus B$ is pre-open.

Remaks 1.3 [4]: 1. Every open (closed) subset of (X, τ) is pre-open (pre-closed) set, but the converse is not true, for example the set of rational numbers is pre-open subset of (\mathbf{R}, τ_u) , where τ_u is an usual topology on \mathbf{R} , but it is not an open set in \mathbf{R} .

2. If G is a subset of X (open) and V is a pre-open set in X then $G \cap V$ is pre-open set.

Definition 1.4 [6]: Let (X, τ) be a topological space, $A \subseteq X$ and let $F = \{V_\alpha : \alpha \in \Omega\}$ be a cover of A is called pre-open cover of A if V_α is a pre-open subset in X for each $\alpha \in \Omega$. Note that every open cover of A is pre-open cover of A .

Definition 1.5 [2]: A topological space (X, τ) is called p -compact if for every pre-open cover of X has a finite subcover.

Remark 1.6: It is known that every p -compact is compact but the converse is not true, for example, let $X = \mathbf{R}$, $\tau = \{\emptyset, X, \{0\}\}$, then X is compact but it is not a pre-compact space.

Definition 1.7 [3]: A subset A of (X, τ) is said to be $G\delta$ -open if for each x in A there is a countable family of open sets $\{U_i\}_{i=1}^\infty$ such that $x \in \bigcap_{i=1}^\infty U_i \subseteq A$. On the other hand A is $G\delta$ -closed if $X \setminus A$ is $G\delta$ -open set.

Note that each open set is $G\delta$ -open and each closed set is $G\delta$ -closed, but the converse are not true.

Definition 1.8 [3]: A function $f : (X, \tau_X) \rightarrow (Y, \tau_Y)$ is called $G\delta$ -closed function if image of each closed subset of X is $G\delta$ -closed subset of Y .

Definition 1.9 [6]: A space (X, τ) is said to be p -T₂-space (p -Hausdorff space) if for each two distinct elements x, y of X there exist two pre-open subset of X , U and V such that $x \in U$, $y \in V$ and $U \cap V = \emptyset$.

Definition 1.10: A subset A of (X, τ) is said to be pre- $G\delta$ -open if for each x in A there is a countable family of pre-open sets $\{U_i\}_{i=1}^\infty$ such that $x \in \bigcap_{i=1}^\infty U_i \subseteq A$. On the other hand A is pre- $G\delta$ -closed if $X \setminus A$ is a pre- $G\delta$ -open. Note that $G\delta$ -closed ($G\delta$ -open) set is a pre- $G\delta$ -closed (pre- $G\delta$ -open) set, but the converse is not true.

Definition 1.11 [6]: A function $f : (X, \tau_X) \rightarrow (Y, \tau_Y)$ is called p^{**} -continuous if $f^{-1}(W)$ is a pre-open set in X for each W is a pre-open set in Y .

Definition 1.12: A topological space (X, τ) is N -space if for every countable intersection of open sets is also open set.

Remark and Examples 1.13: 1. The discrete space (X, D) is N -space.
2. (\mathbf{R}, τ_u) is not N -space.
3. Let (X, τ) be a N -space then every $G\delta$ -open set in X is an open set and $G\delta$ -closed set in X is closed set.

2. Lindelöf spaces and p-Lindelöf spaces

Lindelöf spaces have always played a highly expressive role in topology. They were introduced by Alexandroff and Urysohn back in 1929 and their name is due to Lindelöf's proof in 1903 that from any collection of open sets covering a Euclidean space one can extract a countable subcollection covering the space. The authors in [7] study special kind of Lindelöf space which is called LC-space, but in this section we introduce and study another kind of Lindelöf space which is called p -Lindelöf space. p -Lindelöf space is a generalization of the concept of p -compact space.


Definition 2.1: A topological space (X, τ) is said to be p -Lindelöf space iff every pre-open cover of X has countable subcover.

Remark 2.2: If X is a countable indiscrete space then it is a p -Lindelöf space and if X is an uncountable discrete space it is not p -Lindelöf space.

Proposition 2.3: Every p -Lindelöf space X is a Lindelöf space.

Proof: Let $\{W_\alpha : \alpha \in \Omega\}$ be any open cover for X . Since an open set W_α in X is a pre-open set in X then $\{W_\alpha : \alpha \in \Omega\}$ is a pre-open cover for X , so there is $\alpha_1, \alpha_2, \dots \in \Omega$ such that $X = \bigcup_{i=1}^{\infty} W_{\alpha_i}$ because X is a p -Lindelöf space. Hence $\{W_{\alpha_i}\}$ is a countable open cover for X , therefore X is a Lindelöf space. \square

Remark 2.4: The following diagram shows the relations between compact, p -compact, Lindelöf and p -Lindelöf:



A Lindelöf space need not be p -Lindelöf since an uncountable Lindelöf space is a Lindelöf space, but not p -Lindelöf space.

Proposition 2.5: Every pre-closed subset A of p -Lindelöf space X is a p -Lindelöf space.

Proof: Let $F = \{W_\alpha : \alpha \in \Omega\}$ be any pre-open cover for A . So $F^* = F \cup \{X \setminus A\}$ is a pre-open cover of X , but X is a p -Lindelöf space, hence there is $\alpha_1, \alpha_2, \dots \in \Omega$ such that $X = \bigcup_{i=1}^{\infty} W_{\alpha_i} \cup \{X \setminus A\}$ which implies that $A \subseteq \bigcup_{i=1}^{\infty} W_{\alpha_i}$ i.e. A is a p -Lindelöf space. \square

Remark 2.6: It is known that a continuous function from a compact (p -compact) space into a Hausdorff space is a closed function [2]. However, the continuous function from a Lindelöf (p -Lindelöf) space, need not be closed an example may be found in [4].

Proposition 2.7: The continuous function $f: X \rightarrow Y$ from a Lindelöf X into a T_2 -space Y is a $G\delta$ -closed function.

Proof: Let A be any closed subset of X . In order to prove that $f(A)$ is a $G\delta$ -closed subset of Y , i.e. $Y \setminus f(A)$ is a $G\delta$ -open subset of Y , let $x \in Y \setminus f(A)$ implies that $x \neq y, \forall y \in f(A)$. Since Y is T_2 -space there is U_y, V_y open subsets of Y such that $x \in U_y, y \in V_y$ and $U_y \cap V_y = \emptyset$. Therefore $f(A) \subseteq \bigcup \{V_y : y \in f(A)\}$ so $\{V_y : y \in f(A)\}$ is an open cover for $f(A)$ in Y implies

$$A \subseteq f^{-1}(f(A)) \subseteq f^{-1}(\bigcup \{V_y : y \in f(A)\}) = \bigcup \{f^{-1}(V_y) : y \in f(A)\}$$

But $f^{-1}(V_y)$ is an open subset of X , since f is a continuous function. Thus $\{f^{-1}(V_y) : y \in f(A)\}$ is an open cover for A . By knowing A is a closed subset of Lindelöf space X , then A itself is Lindelöf space which implies there is y_1, y_2, \dots such that $A \subseteq \bigcup_{i=1}^{\infty} f^{-1}(V_{y_i})$, hence $f(A) \subseteq f(\bigcup_{i=1}^{\infty} f^{-1}(V_{y_i})) = \bigcup_{i=1}^{\infty} f(f^{-1}(V_{y_i})) \subseteq \bigcup_{i=1}^{\infty} V_{y_i}$. But for each V_{y_i} there is U_{y_i} such that $U_{y_i} \cap V_{y_i} = \emptyset$, and $x \in U_{y_i}$ for each $i=1, 2, \dots$ implies that $x \in \bigcap_{i=1}^{\infty} U_{y_i} \subseteq Y \setminus f(A)$, hence $Y \setminus f(A)$ is a $G\delta$ -open subset of Y . Therefore, f is a $G\delta$ -closed function. \square

Corollary 2.8: Every continuous function from a p -Lindelöf space into a T_2 -space is a $G\delta$ -closed function.

Proof: Since p -Lindelöf space is a Lindelöf space then use proposition 2.7 we obtain the result. \square

Proposition 2.9: Every Lindelöf subset A of a T_2 -space X is a $G\delta$ -closed set.

Proof: Let $x \in X \setminus A$ and for each $y \in A$ there is U_y, V_y open subsets of X such that $x \in U_y, y \in V_y$ and $U_y \cap V_y = \emptyset$ because X is a T_2 -space. Therefore $\{V_y : y \in A\}$ is an open cover of A , but A is a Lindelöf subset there is y_1, y_2, \dots in A such that $A \subseteq \bigcup_{i=1}^{\infty} V_{y_i}$ and $x \in \bigcap_{i=1}^{\infty} U_{y_i} \subseteq X \setminus A$. Hence $X \setminus A$ is a $G\delta$ -open subset of X i.e. A is a $G\delta$ -closed set. \square

Proposition 2.10: Every p -Lindelöf subset A of a T_2 -space X is a pre- $G\delta$ -closed set.

Proof: Let $x \in X \setminus A$ and for each $y \in A$ there is U_y, V_y pre-open subsets of X such that $x \in U_y, y \in V_y$ and $U_y \cap V_y = \emptyset$ because X is a T_2 -space. Therefore $\{V_y : y \in A\}$ is a pre-open cover of A , but A is a p -Lindelöf subset there is y_1, y_2, \dots in A such that $A \subseteq \bigcup_{i=1}^{\infty} V_{y_i}$ and $x \in \bigcap_{i=1}^{\infty} U_{y_i} \subseteq X \setminus A$. Hence $X \setminus A$ is a pre- $G\delta$ -open subset of X i.e. A is a pre- $G\delta$ -closed set. \square

Proposition 2.11: Every p^{**} -continuous image of a p -Lindelöf space is a p -Lindelöf space.

Proof: Let $f: X \rightarrow Y$ be a p^{**} -continuous function of a p -Lindelöf space X onto a space Y and let $W = \{W_\alpha : \alpha \in \Omega\}$ be any pre-open cover of Y . Then $W^* = \{f^{-1}(W_\alpha) : \alpha \in \Omega\}$ is a pre-open cover of X , because f is a p^{**} -continuous function. Since X is a p -Lindelöf

space W^* has a countable subcover such that $X = \bigcup_{i=1}^{\infty} f(W\alpha_i)$. Since f is onto then $f(f^{-1}(W\alpha_i)) = W\alpha_i$ for each α_i , $i = 1, 2, \dots$, hence $Y = f(X) = \bigcup_{i=1}^{\infty} W\alpha_i$. This implies that Y is a p -Lindelöf space. \square

3. Relative p -Lindelöf and locally p -Lindelöf spaces

We begin with a definition of relative p -Lindelöf and some characterization.

Definition 3.1: A subset A of space X is said to be p -Lindelöf relative to X iff for every cover of A by pre-open sets of X has a countable subcover.

Proposition 3.2: If G is an open subset of space X then G is a p -Lindelöf (as the subspace of X) iff G is a p -Lindelöf relative to X .

Proof: Let G be a p -Lindelöf set in X and let $\{W\alpha : \alpha \in \Omega\}$ be any pre-open cover of G such that $W\alpha$ is a pre-open set of X . By using remark 1.3 (2) we have $G \cap W\alpha$ is a pre-open set in G and in X implies that $\{G \cap W\alpha : \alpha \in \Omega\}$ is a pre-open cover of G . But G is a p -Lindelöf set, thus $G \subseteq \bigcup_{i=1}^{\infty} \{G \cap W\alpha_i\}$ i.e. G is a p -Lindelöf relative to X . Conversely, since G is an open subset of X then it is a p -Lindelöf (see remark 1.3 (1)). \square

Proposition 3.3: Let A and B be two open subsets of X such that $A \subseteq B$ then A is a p -Lindelöf subspace of B if A is a p -Lindelöf subspace of X .

Proof: Let A be a p -Lindelöf relative to B and $\{W\alpha : \alpha \in \Omega\}$ be any pre-open cover of A , where $W\alpha$ is a pre-open set in X . Therefore, $B \cap W\alpha$ is a pre-open set in A . But A is a p -Lindelöf relative to B , hence $A \subseteq \bigcup_{i=1}^{\infty} \{B \cap W\alpha_i\}$ implies $A \subseteq \bigcup_{i=1}^{\infty} \{W\alpha_i\}$ i.e. A is a p -Lindelöf relative to X . Conversely, since any pre-open set in B , where B is an open set in X , is a pre-open set in X (remark 1.3 (2)) then A is a p -Lindelöf relative to B . \square

Definition 3.4 [2]: A space X is said to be locally p -Lindelöf if each element has a neighborhood which is a p -Lindelöf subspace of X .

Remark 3.5: Every p -Lindelöf space is a locally p -Lindelöf space, however the converse may not be true, since an uncountable space is a locally p -Lindelöf but not p -Lindelöf.

Proposition 3.6: A space X is a locally p -Lindelöf if every element of X , there exists an open set which is p -Lindelöf of X .

Proof: Let X be a locally p -Lindelöf, then for each element x in X has a neighborhood which is a p -Lindelöf subspace of X . Thus the element x has a neighborhood which is a p -Lindelöf relative to X (Proposition 3.2). Conversely, let x in X and N_x be an open set to x then N_x is a p -Lindelöf set (Proposition 3.2). Thus N_x is a neighborhood which is a p -Lindelöf subspace of X i.e. X is a locally p -Lindelöf. \square

Proposition 3.7: A space X is a locally p -Lindelöf if for every x in X , there exists an open locally p -Lindelöf subspace A containing x .

Proof: Let X be a locally p -Lindelöf space, then there is a neighborhood N_x such that $x \in N_x$ and N_x is a p -Lindelöf implies that N_x is a locally p -Lindelöf containing x (Proposition 3.6). Conversely, since A is an open locally p -Lindelöf subspace of X

containing x , there exists a neighborhood V of x in A which is a p -Lindelöf subspace of A . Since A is an open set in X , so is V . By using proposition 3.3 V is a p -Lindelöf subspace of X . This shows that X is a locally p -Lindelöf space. \square

REFERENCES :

1. Ganster M., Pre-open sets and resolvable space, Kyungpook Math J. 27 (2) (1987), 135-145.
2. Abd Al-Monsef , M.E. and Noiri , T. , Strongly-compact spaces, Delta J. Sci 8 (1) (1984), 30-46.
3. _____, and El-Deep, S.N., On pre-continuous and weak pre-continuous, Proc. Math - Phys. Soc, Egypt 58 (1982) 47-53.
4. Mustafa, H.J. and Dawood, N.A., C- Lindelöf spaces, to appear.
5. Taheer, N. and Mustafa, H.J., On p -compact spaces, M.Sc. Thesis, Al-Mustansiriyah University, Baghdad (2002).
6. Sharma, J.N., Topology, Meerat (1977).
7. Dontchev J., Ganster M. and Kanibir A., On some generalization of LC-space, Acta Math Univ. Comenianae Vol LXV III 2 (1999) 345-353.